

## Psittacosis - parrot fever

Psittacosis is a type of lung infection caused by the bacterium *Chlamydia psittaci*. This germ is principally carried by birds of the parrot family, including budgerigars, lovebirds and parakeets. Other birds that may harbour the germ include canaries, poultry and pigeons. Both wild and domesticated varieties can carry the bacteria, and some infected birds don't show any sign of illness. The disease is sometimes called 'parrot fever'.

Humans most commonly catch the disease from infected birds by inhaling the bacteria from shed feathers, secretions and droppings. Human-to-human transmission is extremely rare.

Psittacosis can be mild, moderate or severe; some people may have no symptoms. Older people generally experience more severe reactions. The complications of untreated psittacosis include inflammation of the brain or heart. This disease can be readily treated with antibiotics.

### Symptoms of psittacosis

The incubation period for psittacosis is between one week and one month from exposure. Symptoms may include:

- Fever
- Headache
- General malaise
- Muscle aches
- A dry cough
- Shortness of breath.

### People at risk

People who have birds as pets, poultry workers and anyone working in aviaries or pet shops, are most at risk of catching psittacosis. However, even casual contact with an infected bird can result in infection. There have been reports of some people developing psittacosis despite having no recognised contact at all with birds.

### Treatment for psittacosis

Psittacosis is diagnosed with blood tests and chest x-rays. Treatment includes antibiotics. The symptoms usually ease within one day, but the full course of tablets needs to be taken.

Once psittacosis has been diagnosed and if you have pet birds, it is important to test and then if necessary, treat the birds and their environment. Take any sick birds to your veterinarian to allow illness to be investigated. Remember that healthy birds may be harbouring the bacteria as well.

### Preventive measures

Preventive measures include:

- Avoid unnecessary handling of sick birds.
- Avoid breathing in any dust from dried bird droppings, feathers or cage dust.
- Isolate sick birds from the rest of the flock.
- Treat infected birds with appropriate antibiotics for at least one month.
- Clean cages with appropriate disinfectants, since the bacteria can live for several months in shed feathers and droppings.
- Wear masks and gloves while cleaning the cages to prevent infection.

- Clean the cages regularly, using plenty of water to minimise the risk of floating dander.
- Always wash hands thoroughly after tending to birds.

## Infection doesn't confer immunity

Catching psittacosis doesn't confer immunity, which means a person regularly exposed to infected birds can become sick again. There is no vaccine against the disease.

## Where to get help

- Your doctor
- Veterinarian
- Communicable Disease Prevention and Control Unit, Department of Health Victoria Tel. 1300 651 160

## Things to remember

- Psittacosis is a type of lung infection caused by the bacterium *Chlamydia psittaci*.
- *Chlamydia psittaci* is commonly carried by birds of the parrot family including budgerigars, lovebirds and parakeets.
- This disease can be readily treated with antibiotics.

**This page has been produced in consultation with, and approved by:**

Department of Health - Communicable Disease Prevention and Control Unit

Content on this website is provided for education and information purposes only. Information about a therapy, service, product or treatment does not imply endorsement and is not intended to replace advice from your doctor or other registered health professional. Content has been prepared for Victorian residents and wider Australian audiences, and was accurate at the time of publication. Readers should note that, over time, currency and completeness of the information may change. All users are urged to always seek advice from a registered health care professional for diagnosis and answers to their medical questions.

For the latest updates and more information, visit [www.betterhealth.vic.gov.au](http://www.betterhealth.vic.gov.au)

**Copyright** © 1999/2012 State of Victoria. Reproduced from the Better Health Channel ([www.betterhealth.vic.gov.au](http://www.betterhealth.vic.gov.au)) at no cost with permission of the Victorian Minister for Health. Unauthorised reproduction and other uses comprised in the copyright are prohibited without permission.