

IMODIUM(R) Advanced Chewable tablets

Loperamide hydrochloride and simethicone

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about IMODIUM Advanced. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. If you have any concerns about using IMODIUM Advanced ask your doctor or pharmacist.

Keep this leaflet with your medicine.

You may need to read it again.

What IMODIUM Advanced is used for

IMODIUM Advanced is used to treat sudden (acute) diarrhoea.

IMODIUM Advanced also relieves the symptoms of too much gas such as abdominal pain, bloating, cramping and flatulence (wind).

IMODIUM Advanced makes the stools more solid and less frequent. It does this by normalising the muscle contractions of the intestine. This allows the bowel time to absorb water and produce a more solid stool. IMODIUM Advanced also helps the body to get rid of excess gas trapped in the stomach and bowel.

Your doctor may have prescribed IMODIUM Advanced for another reason.

Ask your doctor if you have any questions about why IMODIUM Advanced has been prescribed for you.

Before you take IMODIUM Advanced

When you must not take it:

Do not take IMODIUM Advanced if:

- * you have an allergy to IMODIUM Advanced or any of the ingredients listed at the end of this leaflet (see Product Description).
- * you are constipated, or have to avoid becoming constipated
- * you have severe diarrhoea with blood in the stools or fever
- * you have inflammation of the lower bowels (for example, ulcerative colitis or pseudomembranous colitis as a result of using antibiotics)
- * you are breast-feeding
- * the packaging is torn or show signs of tampering.

Do not use IMODIUM Advanced after the expiry date (month and year) printed on the pack. If you take IMODIUM Advanced after the expiry date it may not work.

Do not give IMODIUM Advanced to children under the age of 12 years.

Before you start to take it:

You must tell your doctor or pharmacist if you:

- * are pregnant or planning to become pregnant
- * are breastfeeding
- * have or have ever had liver disease
- * have kidney disease
- * have AIDS (Acquired Immune Deficiency Syndrome)
- * have glaucoma (pressure in the eye)
- * have bladder problems
- * regularly drink a lot of alcohol.

Taking other medicines:

Tell your doctor or pharmacist if you are taking any other medicines, including medicines you can buy without a prescription from a pharmacy, supermarket or health food shop.

In particular, tell your doctor or pharmacist if you are taking any of the following:

- * sleeping tablets or medicines to treat anxiety and nervousness
- * medicines to treat depression
- * medicines that slow down action of the stomach and intestines.

Taking IMODIUM Advanced

How to take:

IMODIUM Advanced tablet should be chewed, then swallowed. No liquid intake is necessary.

How much to take:

Adults:

The usual adult dose is two IMODIUM Advanced tablets initially, followed by one tablet after each loose unformed stool as needed.

Do not take more than four tablets in one day.

Your doctor may prescribe a different dose for you.

IMODIUM Advanced tablets should be chewed.

Children:

IMODIUM Advanced should not be given to children under 12 years of age.

If you forget to take it:

If you forget to take IMODIUM Advanced after an unformed stool, take it when you remember or after your next unformed stool.

Do not take a double dose to make up for the dose you missed.

If you are not sure what to do, check with your doctor or pharmacist.

If you have taken too much (overdose):

Immediately telephone your doctor or the Poisons Information Centre for advice, or go to Accident and Emergency at your nearest hospital.

Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Poisons Information Centre telephone numbers:

* Australia: 13 11 26

* New Zealand: 0800 POISON or 0800 764 766

Keep these telephone numbers handy in case of an emergency.

If you take too much IMODIUM Advanced you may have any of the following symptoms: sleepiness, nausea and vomiting, constipation, difficulty passing urine, weak breathing, muscle stiffness, and uncoordinated movements.

Children are more sensitive than adults to medicines such as IMODIUM Advanced. An accidental overdose is especially dangerous. Make sure IMODIUM Advanced is kept out of reach of children.

While you are using IMODIUM Advanced

Things you must do:

- * If you do not improve after taking IMODIUM Advanced for 48 hours tell your doctor or pharmacist
- * Drink plenty of fluids, to avoid dehydration
- * Tell your doctor if you become pregnant while taking IMODIUM Advanced
- * Tell your doctor or pharmacist that you are taking IMODIUM Advanced if you are about to start taking a new medicine.

Things you must not do:

- * Do not take more than four tablets in one day
- * Do not use IMODIUM Advanced to treat any other complaint unless your doctor says to
- * Do not give this medicine to anyone else, even if they have the same symptoms as you.

Things to be careful of:

Effects on driving and operating machinery

Tiredness, dizziness or drowsiness can be seen with diarrhoea.

Therefore, it is advisable to use caution when driving or operating machinery.

Side Effects

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some side effects.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor if you experience any of the following and they worry you:

- * nausea or vomiting
- * constipation
- * dry mouth or bad taste
- * headache
- * difficulty passing urine
- * tiredness
- * stomach pain
- * fever
- * dizziness or drowsiness.

These side effects are usually mild. The above list does not represent all side effects that have been reported for IMODIUM Advanced. If you notice side effects not mentioned in this leaflet, please inform your pharmacist or doctor.

STOP taking IMODIUM Advanced and tell your doctor immediately if you experience:

- * asthma, wheezing or shortness of breath
- * swelling of the face, lips or tongue which may cause difficulty in swallowing or breathing

- * sudden or severe blisters, itching, skin rash, hives
- * flaking or peeling of skin
- * cough, chills, fever
- * loss of or depressed level of consciousness

These are serious side effects. You may need urgent medical attention. Serious side effects are rare.

Other side effects not listed above may also occur in some people.

Tell your doctor if you notice anything making you feel unwell when you are taking, or soon after you have finished taking IMODIUM Advanced.

Ask your pharmacist or doctor if you do not understand anything in this list.

After using IMODIUM Advanced

Storage

Keep your IMODIUM Advanced chewable tablets in the pack until it is time to take them.

Keep your IMODIUM Advanced in a cool dry place where the temperature stays below 25 degrees C.

Do not store IMODIUM Advanced, or any other medicine, in the bathroom or near a sink. Do not leave medicines in the car or on windowsills. Heat and dampness can destroy some medicines.

Keep your medicines where children cannot reach them.

A locked cupboard at least one-and-a-half metres (1.5 m) above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking IMODIUM Advanced or your medicine has passed its expiry date, ask your pharmacist what to do with any medicine that is left over.

Product Description

What IMODIUM Advanced looks like:

IMODIUM Advanced chewable tablets are white tablets embossed with IMO.

Each pack contains 6 or 12 tablets.

Ingredients

Each IMODIUM Advanced chewable tablet contains 2mg of loperamide hydrochloride and 125 mg of simethicone as the active ingredients.

The chewable tablets also contain confectioners sugar (containing sucrose and maize starch), microcrystalline cellulose, methacrylic acid copolymer, cellulose acetate, sorbitol, dextrates, saccharin sodium, stearic acid, calcium phosphates and Flavorburst(R) Natural and Artificial Vanilla Flavor.

The chewable tablets do not contain lactose or gluten.

Sponsor

Johnson & Johnson Pacific
45 Jones Street
Ultimo NSW 2007
Australia

For further enquiries call our Toll Free Info Line
Aus 1800 029 979
NZ 0800 446 147

Australian Register Number: AUST R 66848

This leaflet was prepared in March 2009.