

Bursitis

Bursitis is inflammation of a bursa. Bursae are small fluid filled sacs located between two adjoining structures, usually muscles, tendons and bones. Bursae decrease friction and assist movement of the tendon over the bony surface. Bursitis does not cause joint deformity but can cause significant pain and restrict movement.

Bursae are located outside the joint itself. Since both tendons and bursae are located near joints, inflammation in these soft tissues may be mistaken as arthritis. However, arthritis involves inflammation within joints, whereas bursitis involves inflammation outside the joint.

Bursitis is most common in the shoulder, elbow, hip and knee

The body contains many bursae but the shoulder, elbow, hip, knee and heel are the most common sites of bursitis. Common causes include injury, repeated pressure and overuse. Certain disorders (such as rheumatoid arthritis, gout and diabetes) can also contribute to the development of bursitis.

Symptoms of bursitis

The symptoms of bursitis may include:

- Localised pain
- Swelling
- A warm feeling in or around the affected area
- Increased pain at night
- Pain worsened by movement
- Stiffness
- Reddening of the skin.

Overuse injury

A common cause of bursitis is overuse of the particular body part, especially if that activity is performed awkwardly or with considerable pressure. Examples of work-related activities that may trigger bursitis include production line packing and typing. Sports that can cause bursitis include jogging, tennis and squash.

Other soft tissue injuries that can be classified as overuse injuries include:

- **Carpal tunnel syndrome** – a painful disorder of the hand leading to increased pressure on the main nerve that runs through the wrist.
- **Ganglion** – a cyst on a tendon or joint capsule.
- **Tennis elbow** – inflammation of tendons surrounding the elbow joint.
- **Tenosynovitis** – inflammation of the tendons (the tough connective tissue that anchors muscle to bone).

Diagnosis of bursitis

Bursitis is generally detected as a tender, warm swelling at the site of a bursa. A diagnosis may also include investigating and ruling out any other possible causes. Tests may include:

- Physical examination
- Medical history

- X-rays, to rule out the possibility of any other condition
- Ultrasound
- Drawing off fluid from the bursa to rule out the possibility of infection.

Treatment for bursitis

Treatment will depend on the cause of the bursitis. Treatment aims to alleviate the symptoms as much as possible while the healing process takes place. Options may include pain-killing drugs, cold packs, gentle mobilising exercises and rest. Anti-inflammatory medications or injections of corticosteroids may be used in cases of severe pain.

If infection is present, there is usually warmth, redness, pain and swelling in the affected areas. Treatment with an appropriate antibiotic is necessary. If the bursitis was triggered by overuse, it is important to avoid the particular activity.

Correct posture and joint protection are useful and braces or splints can decrease the stress on the areas and support good alignment. After an acute attack, it is important to consider how recurrences can be prevented.

Long-term management of bursitis

Your doctor or physiotherapist can offer suggestions and strategies to reduce your risk of developing bursitis again. General suggestions include:

- **Work-related bursitis** – use ergonomically designed furniture and equipment, take regular breaks, do simple stretching exercises for 10 minutes every hour and keep benches at waist height so that the shoulders can relax.
- **Sport-related bursitis** – warm up thoroughly by stretching and gently going through the motions of your chosen sport. Make sure you use good form and regularly practice strengthening and conditioning exercises that compliment your particular sport. Cool down thoroughly with gentle, sustained stretches. Make sure footwear and equipment are appropriate for you.

Where to get help

- Your doctor
- Physiotherapist
- Arthritis Victoria Tel. (03) 8531 8000
- Australian Physiotherapy Association Tel. (03) 9092 0888

Things to remember

- Bursitis is inflammation of a bursa.
- Bursae are small sacs that contain fluid to lubricate moving parts, such as joints, muscles and tendons.
- Common causes of bursitis include injury and overuse.
- Treatment includes rest, ice packs, gentle mobilisation exercises and avoiding the particular activity that prompted the condition.

This page has been produced in consultation with, and approved by:

Arthritis Victoria incorporating Osteoporosis Victoria

Content on this website is provided for education and information purposes only. Information about a therapy, service, product or treatment does not imply endorsement and is not intended to replace advice from your doctor or other registered health professional. Content has been prepared for Victorian residents and wider Australian audiences, and was accurate at the time of publication. Readers should note that, over time, currency and completeness of the information may change. All users are urged to always seek advice from a registered health care professional for diagnosis and answers to their medical questions.

For the latest updates and more information, visit www.betterhealth.vic.gov.au

Copyright © 1999/2012 State of Victoria. Reproduced from the Better Health Channel (www.betterhealth.vic.gov.au) at no cost with permission of the Victorian Minister for Health. Unauthorised reproduction and other uses comprised in the copyright are prohibited without permission.