

## Gastroenteritis - giardiasis

Giardiasis is a form of gastroenteritis, also known as gastro. It is caused by a parasite called *Giardia Lamblia*, which lives in the bowel. Giardiasis can affect anyone but is more common in infants, children and adults aged 20 to 40 years. It is easily spread in childcare centres where there are children who are not toilet trained. *Giardia* parasites are also found in wild, pet and farm animals. Untreated water that comes directly from lakes and rivers may also contain *Giardia* parasites.

### Symptoms that may develop

Symptoms usually take an average of seven to 10 days to develop, but may take three weeks or occasionally longer. Symptoms may include:

- Acute or chronic diarrhoea
- Weight loss
- Fatigue
- Abdominal cramps
- The excretion of large amounts of fat in faeces (steatorrhoea) due to malabsorption of fat in the digestive system.

### Many people do not develop symptoms

Most people who become infected with *Giardia* parasites do not develop symptoms but can still spread the infection to others.

### Poor hygiene spreads the infection

People with *Giardia* parasites in their faeces can infect others if they do not wash their hands properly after going to the toilet. Contaminated hands can then spread the parasites to food that may be eaten by other people or to surfaces that may be touched by other people. Hands can also become contaminated when changing the nappy of an infected infant or handling infected animals.

Giardiasis can be spread by contaminated drinking water. It can also be spread by faecal contamination of public water supplies or recreational swimming areas, such as pools and spas. People who have had giardiasis should not use swimming pools until at least one week after their symptoms have ceased.

### Tell your doctor

If you think that you or your child has giardiasis contact your doctor. Your doctor will arrange for a stool (faeces) sample to be tested. If the results show giardiasis, the doctor will provide appropriate advice and treatment. Your doctor will also notify the Department of Health Victoria.

### How to prevent the spread of giardiasis

The spread of giardiasis can be reduced if you:

- Have the illness diagnosed and treated by your doctor.
- Keep children home from school or child care until symptoms have stopped.
- Stay home from work until symptoms have stopped if you are a food handler, health care worker or a childcare worker.
- Wash hands thoroughly after going to the toilet, changing nappies or handling animals.
- Clean bathrooms and toilets thoroughly.
- Do not drink untreated water.
- Do not swim in public pools until at least one week after symptoms have stopped.
- Do not prepare or handle food that will be eaten by other people.
- Do not share any towel or face washer with a person who has giardiasis.

### Where to get help

- Your doctor
- Your local council health department
- Communicable Disease Prevention and Control Unit, Department of Health Victoria Tel. 1300 651 160

### **Things to remember**

- Giardiasis is spread by poor hygiene practises.
- Many people infected with *Giardia* parasites will not have symptoms but can still pass on the infection.
- It is important that you contact your doctor if you think that you or your child has giardiasis.

**This page has been produced in consultation with, and approved by:**

Department of Health - Communicable Disease Prevention and Control Unit

**Copyright** © 1999/2009 State of Victoria. Reproduced from the Better Health Channel ([www.betterhealth.vic.gov.au](http://www.betterhealth.vic.gov.au)) at no cost with permission of the Victorian Minister for Health. Unauthorised reproduction and other uses comprised in the copyright are prohibited without permission.

- This Better Health Channel fact sheet has passed through a rigorous approval process. For the latest updates and more information visit [www.betterhealth.vic.gov.au](http://www.betterhealth.vic.gov.au).