

Erectile dysfunction treatments

A range of treatments is available for erectile dysfunction (ED), or impotence. These include oral medications (tablets) such as Viagra, Cialis or Levitra or injections into the penis. Other treatments include vacuum devices, counselling or implants (penile prosthesis) or – rarely – vascular surgery.

Most cases of erectile dysfunction are caused by physical illness with only one in ten cases of persistent ED caused by psychological problems.

When to see a doctor

Occasional erectile dysfunction is normal and may be a result of too much alcohol, stress or anxiety, or a lack of sleep. However, ongoing impotence can sometimes be a symptom of a serious physical illness. Some of these illnesses include heart and blood vessel diseases (atherosclerosis), diabetes and multiple sclerosis.

Ongoing erection problems should always be fully investigated by a doctor to check for any underlying medical cause that may also require treatment.

Treatments are varied

Sometimes direct treatment of the cause may restore erectile function, for example, through testosterone replacement, resolving substance abuse problems or withdrawing medications. Mostly, treatment aimed at directly restoring erectile function is necessary.

There are many treatments aimed at restoring erectile function. Treatments can be grouped according to their level of invasiveness.

- **Non-invasive treatments** – include oral medications (tablets), counselling and external devices such as vacuum pumps.
- **Minimally invasive treatments** – include penile injections.
- **Surgical treatments** – include penile prosthesis (implants) and vascular surgery.

It is usual to start with the least invasive treatment, such as tablets, and to recommend more complicated injection treatments or surgery if the tablets fail or are inappropriate.

Medications

Oral medication is effective in about 70 per cent of men with erectile dysfunction. This is usually taken in the form of tablets – brand names include Viagra, Cialis and Levitra. These drugs, known as PDE5 inhibitors, inhibit a particular enzyme in the penis, enhancing the erectile response to normal sexual stimulation.

It is usually recommended that tablets be taken at least one hour before sexual contact is planned. Fatty meals and large amounts of alcohol can interfere with the body's absorption of some of these drugs. Side effects of these medications may include facial flushing, nasal congestion, headaches and indigestion. Although these drugs act in a similar way, there are important differences between them – for example, the period of time for which they are active.

Your doctor will advise you about which is the most appropriate for you. Oral medication treatment may not be suitable for all men.

Patients taking nitrate-containing heart medication should always consult their doctor, as erectile dysfunction medications may interact with these medications.

Psychotherapy and counselling

Psychosocial issues may be a single cause of erectile dysfunction or may combine with a physical cause. Psychotherapy and counselling can help if the underlying cause of erectile dysfunction is psychological. Counselling can also benefit the man who has lost sexual confidence, even though his erectile dysfunction is caused by physical factors.

Counselling may be provided by your doctor, a psychologist or a psychiatrist. The man's partner may also be involved.

Vacuum and other devices

A vacuum device is a pump placed over the penis to create blood flow. It draws blood into specific tissues of the penis (the two corpora cavernosa), mimicking a normal erection.

A rubber ring can sometimes be used around the base of the penis to help make it rigid enough for intercourse. This is used when a man is able to get an erection but has difficulty maintaining it.

Devices such as vacuums and rings are not always successful. These devices need to be properly explained and appropriately fitted and are best used after consultation with a doctor.

Injections

Some medications can increase blood flow when they are injected into the penis. A penile injection can create an erection within five to 10 minutes and last for up to one hour. Men must be taught how to use the penile injections properly and should have several supervised injections in a doctor's surgery.

One possible side effect is a painful erection lasting for hours, known as priapism. This occurs if the medication dose is too high. If an erection lasts for four hours or longer, the penis may be permanently damaged. Immediate medical attention should be sought in this situation.

Testosterone replacement therapy

Testosterone replacement therapy (TRT) is only suitable for men whose levels of testosterone are low. Prior to treatment the cause of low testosterone needs to be investigated.

Penile prosthetic implants

A penile prosthesis is a device that is surgically implanted into the penis. It can enable erections through a mechanism of squeezing on a specific part of the device. Penile prosthetic implants are expensive and are generally used only as a last resort if other treatments have not worked.

Vascular surgery

Vascular surgery targets the veins and arteries that supply blood to the penis. It is a major surgical procedure and is rarely used. Surgery may be considered for men who have vascular damage due to trauma in the area around the penis – for example, after a car accident or fall.

Other products

Many other products claim to treat or even cure erectile dysfunction. These products are often expensive and usually have not been adequately tested for either their side effects or results. Treat such products with caution and use only under the supervision of a doctor.

Where to get help

- Your doctor
- Melbourne Sexual Health Centre Tel. (03) 9341 6200 or 1800 032 017 or TTY (for the hearing impaired) (03) 9347 8619 www.mshc.org.au
- Family Planning Victoria Tel. (03) 9257 0100 or 1800 013 952 www.fpv.org.au

Things to remember

- Most cases of erectile dysfunction are caused by physical factors.
- Treatments include oral medications, penile injections, vacuum devices, counselling and implants.
- Ongoing erection problems may have a medical cause that needs specific treatment and should be fully investigated by a doctor.

This page has been produced in consultation with, and approved by:

Andrology Australia

Content on this website is provided for education and information purposes only. Information about a therapy, service, product or treatment does not imply endorsement and is not intended to replace advice from your doctor or other registered health professional. Content has been prepared for Victorian residents and wider Australian audiences, and was accurate at the time of publication. Readers should note that, over time, currency and completeness of the information may change. All users are urged to always seek advice from a registered health care professional for diagnosis and answers to their medical questions.

For the latest updates and more information, visit www.betterhealth.vic.gov.au

Copyright © 1999/2011 State of Victoria. Reproduced from the Better Health Channel (www.betterhealth.vic.gov.au) at no cost with permission of the Victorian Minister for Health. Unauthorised reproduction and other uses comprised in the copyright are prohibited without permission.