
Bowel cancer screening

Summary

- Bowel cancer is the second most common cause of cancer in Australia.
 - More than ninety per cent of bowel cancers can be successfully treated if they are found early.
 - For most people, screening for bowel cancer involves a simple, at-home screening test.
 - The National Bowel Cancer Screening Program (NBCSP) sends free tests in the mail to people aged 50 – 74, every two years.
 - The NBCSP bowel cancer screening test kit looks for traces of blood in the bowel movement which could be a sign of bowel cancer.
 - Participation in the NBCSP is the most effective way people aged 50–74 can prevent bowel cancer.
-

Bowel cancer is the second most common cause of cancer in Australia, with around 100 people dying from the disease each week. Once you reach 50 years of age, your risk of bowel cancer increases.

Screening for bowel cancer in people who do not have any symptoms and are between 50 and 74 years helps to detect cancer early. More than 90 per cent of bowel cancers can be successfully treated if they are detected early.

The **National Bowel Cancer Screening Program (NBCSP)** sends free bowel cancer screening tests in the mail to eligible people between the ages of 50 and 74 years. From 2020, all eligible people between 50 and 74 years will receive free tests every two years.

National Bowel Cancer Screening Program

The National Bowel Cancer Screening Program sends screening tests in the mail to eligible people aged 50 to 74 years. In 2019, the free at home test will be sent to all Australians aged 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72 and 74. From 2020, all eligible Australians aged 50-74 will be invited to screen every two years. This could save up to 500 lives annually and significantly reduce the burden of bowel cancer on Australians and their families.

The National Health and Medical Research Council recommends screening every two years for people aged 50 to 74 years who do not have symptoms or a family history of bowel cancer. The screening test offered is called an immunochemical faecal occult blood test (FOBT).

Bowel cancer screening test

Bowel cancer can grow for years in the wall of the bowel before it spreads to other parts of the body. During this time it may not cause any symptoms.

A growth known as a **polyp** can turn into bowel cancer. Polyps and early-stage bowel cancer can release tiny amounts of blood that cannot be seen with the human eye. Blood in your bowel movement could be an indication of polyps or bowel cancer, but could also be caused by haemorrhoids or other conditions.

Regular FOBT screening can help detect changes in the bowel that could lead to a diagnosis of bowel cancer. An FOBT can detect tiny amounts of blood in bowel movements and pick up polyps or early-stage cancer. If blood is detected, you and your doctor will both receive the results and it is important to visit your doctor for follow-up, which is likely to be a referral for a **colonoscopy**.

The test results can take a few days to come back. It is very natural to feel anxious while waiting to get your results. It can help to talk to a close friend or relative about how you are feeling. You can also contact the **Cancer**

Council on **13 11 20** and speak with a cancer nurse.

Negative result for the bowel cancer screening test

A negative result from the FOBT means that blood was not detected in your bowel movement. This does not mean you will never get bowel cancer or that you definitely do not have cancer.

If you receive a negative result but notice unusual changes in your bowel habits or bleeding from the bowel, visit your doctor. If you do not have symptoms of **bowel cancer**, you should continue to complete the NBCSP test when you receive it in the mail every two years.

Positive result for the bowel cancer screening test

A positive result from the FOBT means that blood was detected in your bowel movement. Contact your doctor immediately to discuss the result and obtain a referral for further investigation, which is likely to be a **colonoscopy**.

The presence of blood in your bowel movement may be due to conditions other than cancer, such as polyps, haemorrhoids or inflammation of the bowel. Even so, you will need further investigation to identify the cause of the bleeding. Having a colonoscopy may enable:

- the cause of the bleeding to be found
- any polyps to be removed
- a diagnosis of bowel cancer to be confirmed (if applicable).

Where to get help

- Your **GP (doctor)**
- Local community health centre
- **National Bowel Cancer Screening Program** Tel. **1800 118 868**
- **Cancer Council**. Tel. **13 11 20**

This page has been produced in consultation with and approved by:

Cancer Council Victoria

Content on this website is provided for information purposes only. Information about a therapy, service, product or treatment does not in any way endorse or support such therapy, service, product or treatment and is not intended to replace advice from your doctor or other registered health professional. The information and materials contained on this website are not intended to constitute a comprehensive guide concerning all aspects of the therapy, product or treatment described on the website. All users are urged to always seek advice from a registered health care professional for diagnosis and answers to their medical questions and to ascertain whether the particular therapy, service, product or treatment described on the website is suitable in their circumstances. The State of Victoria and the Department of Health & Human Services shall not bear any liability for reliance by any user on the materials contained on this website.

For the latest updates and more information, visit www.betterhealth.vic.gov.au

Copyright © 1999/2020 State of Victoria. Reproduced from the Better Health Channel (www.betterhealth.vic.gov.au) at no cost with permission of the Victorian Minister for Health. Unauthorised reproduction and other uses comprised in the copyright are prohibited without permission.