
Fainting

Summary

- Common causes of fainting include heat, pain, distress, the sight of blood, or anxiety and hyperventilating.
 - Lying the person down will often improve the person's condition.
 - Frequent fainting spells need to be medically investigated to check for underlying causes.
-

Fainting is a brief episode of unconsciousness caused by a sudden drop in blood pressure. The most likely cause of this sudden drop will either be some change in the blood vessels or the heartbeat itself.

Blood vessels continually adjust their width to ensure a constant blood pressure. For instance, the vessels constrict (tighten) when we stand up to counteract the effects of gravity. Temporary low blood pressure can be caused by various events that prompt blood vessels to dilate (expand), including extreme heat, emotional distress or pain. The lack of blood to the brain causes loss of consciousness.

Most fainting will pass quickly and won't be serious. Usually, a fainting episode will only last a few seconds, although it will make the person feel unwell and recovery may take several minutes. If a person doesn't recover quickly, always seek urgent medical attention.

Symptoms of fainting

The symptoms of a faint include:

- Dizziness
- Light-headedness
- A pale face
- Perspiration
- Heightened anxiety and restlessness
- Nausea
- Collapse
- Unconsciousness, for a few seconds
- Full recovery after a few minutes.

Occasionally, a collapse may be caused by a more serious event such as a stroke or a disturbance in the normal heart rhythm. A faint might be telling you something is wrong and further examination is sometimes important.

If a person complains of breathlessness, chest pains or heart palpitations, or if the pulse is faster or slower than expected, the person should see a doctor. Similarly, slurred speech, facial droop or weakness in any limbs are signs of a serious problem.

Causes of a drop in blood pressure

A temporary drop in blood pressure can be caused by different factors, including:

- Prolonged standing
 - Extreme heat, which pushes blood away from the main circulatory system and into the vessels of the skin
 - Emotional distress
 - Severe pain
 - The sight of blood
 - The sight of a hypodermic needle
-

- Other events that a person may find distressing.

What to do if you feel faint

If possible, lie down and elevate the feet. This may prevent a loss of consciousness. Fresh air can also help, especially if you are feeling hot. If it is not possible to lie down, put your head down as low as possible.

If you do faint, remain lying down for ten minutes. Sit up slowly when you need to get up.

First aid and fainting

First aid treatment for a person who has fainted includes:

- Help the person lie down. A person who has fainted in a chair should be helped to the ground.
- If the person is unconscious, roll them on their side. Check they are breathing and that they have a pulse.
- If possible, elevate the person's feet above the height of their head.
- If the fainting episode was brought on by heat, remove or loosen clothes, and try to cool the person down by wiping them with a wet cloth or fanning them.
- Assess the person for any potential injuries if they have fallen.
- In an emergency, always call triple zero (000) for an ambulance if the person has not regained consciousness within a few seconds or recovered in a few minutes.

Hypotension and fainting

Low blood pressure (hypotension) is a condition characterised by blood pressure that is lower than normal or usual for the person.

Hypotension can be caused by a variety of factors including heart disease and abnormal heart rhythms, some infections, dehydration and medications for high blood pressure or certain heart conditions. Low blood pressure can also be caused by a rare disorder of the adrenal glands called Addison's disease. Frequent fainting spells or sensations of light-headedness need to be medically investigated to check for underlying causes.

Orthostatic hypotension

Blood vessels respond to gravity by constricting (tightening). This increases or maintains blood pressure when we stand up from a sitting or lying position.

Orthostatic hypotension means that the blood vessels don't adjust to a standing position, but instead allow the blood pressure to drop, which can trigger a fainting episode. For this reason, some people, particularly the elderly or those on blood pressure medication, should stand up from sitting or lying in bed slowly. This helps prevent fainting after sudden changes in position.

Causes of orthostatic hypotension include:

- Nervous system diseases, such as neuropathy
- Prolonged bed rest
- Dehydration
- Irregular heartbeat (heart arrhythmia)
- Changes in blood pressure medication.

Where to get help

- Your doctor
- In an emergency, always call triple zero (000).

Things to remember

- Common causes of fainting include heat, pain, distress, the sight of blood, or anxiety and hyperventilating.
- Lying the person down will often improve the person's condition.
- Frequent fainting spells need to be medically investigated to check for underlying causes.

This page has been produced in consultation with and approved by:

Ambulance Victoria

Content on this website is provided for information purposes only. Information about a therapy, service, product or treatment does not in any way endorse or support such therapy, service, product or treatment and is not intended to replace advice from your doctor or other registered health professional. The information and materials contained on this website are not intended to constitute a comprehensive guide concerning all aspects of the therapy, product or treatment described on the website. All users are urged to always seek advice from a registered health care professional for diagnosis and answers to their medical questions and to ascertain whether the particular therapy, service, product or treatment described on the website is suitable in their circumstances. The State of Victoria and the Department of Health & Human Services shall not bear any liability for reliance by any user on the materials contained on this website.

For the latest updates and more information, visit www.betterhealth.vic.gov.au

Copyright © 1999/2020 State of Victoria. Reproduced from the Better Health Channel (www.betterhealth.vic.gov.au) at no cost with permission of the Victorian Minister for Health. Unauthorised reproduction and other uses comprised in the copyright are prohibited without permission.