

---

## Fever - febrile convulsions

---

### Summary

- A febrile convulsion is a fit or seizure that occurs in children when they have a high fever.
  - A febrile convulsion is not epilepsy and a short-lived fit will not cause brain damage – even a long fit almost never causes harm.
  - Call an ambulance if the fit lasts longer than five minutes, as drugs may be needed to stop the fit.
- 

A febrile convulsion is a fit or seizure that occurs in children when they have a high fever. This can happen in children aged six months to five years. The fit can last a few seconds or up to 15 minutes and is followed by drowsiness. Most fits last less than two to three minutes.

One in every 20 children will have one or more febrile convulsion. A febrile convulsion is not epilepsy and does not cause brain damage. Around 30 per cent of babies and children who have had one febrile convulsion will have another. There is no way to predict who will be affected or when this will happen.

### Symptoms of febrile convulsions

The symptoms include:

- loss of consciousness (black out)
- twitching or jerking of arms and legs
- breathing difficulty
- foaming at the mouth
- going pale or bluish in skin colour
- eye rolling, so only the whites of their eyes are visible
- your child may take 10 to 15 minutes to wake up properly afterwards. They may be irritable during this time and appear not to recognise you.

### Reassurance for parents and febrile convulsions

The signs and symptoms of a febrile convulsion can be very frightening to parents. Important things to remember include:

- Children suffer no pain or discomfort during a fit.
- A febrile convulsion is not epilepsy. No regular drugs are needed.
- A short-lived fit will not cause brain damage. Even a long fit almost never causes harm. Children who have had a febrile convulsion grow up healthy.
- If you have concerns or questions, contact your local doctor. In an emergency, take your child to the nearest hospital emergency department.
- There is a medication called Midazolam that is sometimes recommended for children who have a history of febrile convulsions lasting longer than five minutes. Most children do not require this medication. If you would like more information about this treatment, you should talk with your doctor.

### Causes of febrile convulsions

Febrile convulsions only happen when there is a sudden rise in body temperature. The fever is usually due to a viral illness or, sometimes, a bacterial infection. The growing brain of a child is more sensitive to fever than an adult brain. Febrile convulsions tend to run in families, although the reason for this is unknown.

### Treatment for a fever

---

Fever is a normal response to infection and is usually harmless. If your child has a fever, suggestions include:

- Keep them cool by not overdressing them or having their room too hot.
- Give them plenty to drink. It is best to give small, frequent drinks of water.
- Give liquid paracetamol if your child has pain or is miserable. Check the label for how much to give and how often. Paracetamol does not protect against febrile convulsions.

### **First aid for febrile convulsions**

If your child experiences a fit, suggestions include:

- Try to stay calm and don't panic.
- Make sure your child is safe by placing them on the floor. Remove any object that they could knock themselves against.
- Don't force anything into your child's mouth.
- Don't shake or slap your child.
- Don't restrain your child.
- Once the convulsion has stopped, roll your child onto their side, also known as the recovery position. If there is food in their mouth, turn their head to the side, and do not try to remove it.
- Note the times that the fit started and stopped to tell the doctor.
- Have your child checked by your local doctor or nearest hospital emergency department as soon as possible after the fit stops.
- Call an ambulance if the fit lasts longer than five minutes, as medications may be needed to stop the fit.

### **Where to get help**

- Your doctor
- Emergency department of your nearest hospital
- Always call an ambulance in an emergency Tel. 000
- Maternal and Child Health Line Tel. 13 22 29 (24 hours)
- NURSE-ON-CALL Tel. 1300 606 024 – for expert health information and advice (24 hours, 7 days)
- The Royal Children's Hospital Tel. (03) 9345 5522

### **Things to remember**

- A febrile convulsion is a fit or seizure that occurs in children when they have a high fever.
- A febrile convulsion is not epilepsy and a short-lived fit will not cause brain damage – even a long fit almost never causes harm.
- Call an ambulance if the fit lasts longer than five minutes, as drugs may be needed to stop the fit.

**This page has been produced in consultation with and approved by:**

The Children's Hospital at Westmead

Content on this website is provided for information purposes only. Information about a therapy, service, product or treatment does not in any way endorse or support such therapy, service, product or treatment and is not intended to replace advice from your doctor or other registered health professional. The information and materials contained on this website are not intended to constitute a comprehensive guide concerning all aspects of the therapy, product or treatment described on the website. All users are urged to always seek advice from a registered health care professional for diagnosis and answers to their medical questions and to ascertain whether the particular therapy, service, product or treatment described on the website is suitable in their circumstances. The State of Victoria and the Department of Health & Human Services shall not bear any liability for reliance by any user on the materials contained on this website.

For the latest updates and more information, visit [www.betterhealth.vic.gov.au](http://www.betterhealth.vic.gov.au)

**Copyright © 1999/2019** State of Victoria. Reproduced from the Better Health Channel ([www.betterhealth.vic.gov.au](http://www.betterhealth.vic.gov.au)) at no cost with permission of the Victorian Minister for Health. Unauthorised reproduction and other uses comprised in the copyright are prohibited without permission.