
Leukaemia

Summary

- Leukaemia is a cancer of blood-forming cells, which causes large numbers of white blood cells to be made.
 - A lot of leukaemia can be cured or kept under control for many years.
 - There are four main types of leukaemia.
 - The main treatment is with chemotherapy.
-

Leukaemia is a cancer of the blood-forming cells, which are made in the bone marrow. Leukaemia causes large numbers of white blood cells to be produced. This can crowd the bone marrow and may affect the production of normal cells. A person with leukaemia is more likely to pick up infections and have trouble getting rid of them. About 790 Victorians develop leukaemia every year.

There are four main types of leukaemia. In many cases, leukaemia can be cured or kept under control for many years. Chemotherapy is the main treatment.

Blood cells and leukaemia

Blood is mostly made up of plasma, which is a clear fluid. Floating in this fluid are three types of cells, which are:

- red blood cells – that carry oxygen around the body
- white blood cells – that fight disease and infection
- platelets – that help to stop bleeding when it starts.

Blood cells are made in the bone marrow, which is the spongy part in the centre of the bone.

When the body is healthy, the numbers of red cells, white cells and platelets in the blood are kept in balance. In people with leukaemia, the bone marrow makes too many abnormal white blood cells. These abnormal cells are unable to carry out their normal function and can crowd the bone marrow. This means there is not enough room in the bone marrow for making normal white blood cells, red blood cells and platelets.

White blood cells normally help to fight infection. In a person with leukaemia, a large number of these cells are abnormal. If a person has too many abnormal white blood cells, they are more likely to pick up and have trouble getting rid of infections.

Causes of leukaemia

The cause of most cases of leukaemia is unknown. There are several risk factors that may increase your chance of developing this disease.

The main ones are:

- exposure to intense levels of radiation – no cases of leukaemia have been linked to radiation from x-rays and CT scans
- exposure to the chemical benzene – happens in a work environment over long periods of time
- cigarette smoking – this can increase your risk of some types of leukaemia
- genetic factors – certain inherited conditions can increase your risk of developing some types of leukaemia
- family history, age and gender – these may all have an effect on your risk of developing certain types of leukaemia (for example, males are at a higher risk of developing leukaemia than females).

Research into further possible causes of leukaemia is ongoing.

Types of leukaemia

There are different types of leukaemia and the treatment for each is different. The condition may:

- develop suddenly (over days or weeks) – this is called acute leukaemia
- develop over months or years – this is called chronic leukaemia
- involve either the myeloid or lymphoid cells – the condition is named for the type of blood cell affected.

The four main types of leukaemia are:

- acute lymphocytic leukaemia (ALL)
- chronic lymphocytic leukaemia (CLL)
- acute myeloid leukaemia (AML)
- chronic myeloid leukaemia (CML).

Acute lymphocytic leukaemia (ALL)

ALL affects lymphocytes (a type of white blood cell, whose main function is to protect the body from infection) so they cannot work properly. This may cause a serious infection. It also causes many abnormal lymphocytes to be made, which crowd out the normal red blood cells and platelets. ALL is the most common type in children. It is rare in adults.

Chronic lymphocytic leukaemia (CLL)

CLL also affects the lymphocytes, but develops more slowly than ALL. Most people with CLL are not even aware that they have the disease and may only be diagnosed during an examination for another medical problem. CLL affects adults and does not occur in children.

Acute myeloid leukaemia (AML)

AML mainly affects the myeloid cells known as granulocytes. These cells search for infectious invaders in the blood and nearby tissue. In AML, too many young myeloid cells are produced and there are not enough mature myeloid cells. The young myeloid cells can block blood vessels. AML mainly affects adults, but can occur in children and adolescents.

Chronic myeloid leukaemia (CML)

Too many mature myeloid cells are present in CML and they do not function properly. CML is also different from the more common types of leukaemia because it has two stages. In the first stage, abnormal cells slowly multiply, and then in the second stage, it quickly changes and becomes exactly like AML.

This leukaemia can occur at any age, but is uncommon under the age of 20 years.

Symptoms of leukaemia

The symptoms of leukaemia may vary slightly depending on the type and stage of leukaemia you have.

Symptoms for acute leukaemia usually happen quite suddenly. They include:

- weakness, tiredness and looking 'washed out'
- regular infections and high temperatures
- bleeding easily
- bleeding from the gums and nose
- low red blood cells (anaemia) causing breathlessness and tiredness (more common in later stages of CLL)
- bruising easily with no obvious cause
- pain in the bones and joints
- weight loss
- swollen lymph glands
- abdominal discomfort due to an enlarged spleen.

People with chronic leukaemia will also have many of these symptoms, but they tend to develop over months or

years.

Most people who have any of these symptoms will not have leukaemia. They are more likely to be suffering from a much less serious problem, such as an infection. However, it is always important to see your doctor if you have symptoms that persist.

Diagnosis of leukaemia

Tests to help diagnose leukaemia may include:

- Blood tests – these check for large numbers of abnormal white blood cells.
- Bone marrow biopsy – a small amount of bone marrow is taken from the hip bone using a needle. The samples are then examined under a microscope. This test can help work out which type of leukaemia is present.
- Lymph node biopsy – a small amount of tissue is removed with a needle from a swollen lymph node and examined under a microscope. This is more commonly used for lymphoma than leukaemia.
- Lumbar puncture – fluid is removed with a needle from a space between the bones in the back. The fluid is looked at under a microscope.

If these tests show that you do have leukaemia, you may also need to have further blood and bone marrow tests, as well as some scans (CT, ultrasound or MRI).

Test results can take a few days to come back. It is very natural to feel anxious while waiting to get your results. It can help to talk to a close friend or relative about how you are feeling. You could also contact the Cancer Information and Support Service on 13 11 20 and speak with a cancer nurse.

Treatment of leukaemia

Most children and many adults with acute leukaemia can expect to be cured. For most people, chronic leukaemia can be successfully managed for long periods of time.

Treatment depends on the type of leukaemia, but may include:

- **Chemotherapy** – medication to treat cancer by destroying or slowing the growth of fast-growing cancer cells. This is usually given intravenously, but may sometimes be given as tablets. For some people, chemotherapy for acute leukaemia will mean spending several weeks in hospital. Others may be able to stay at home, but will need regular hospital check-ups and further treatments.
- **Stem cell transplant** (stem cells are immature white blood cells) – a transplant allows you to have much higher doses of chemotherapy than usual. This helps improve the chances of curing some types of leukaemia or of prolonging remission. The high doses of chemotherapy, sometimes given with total body irradiation (TBI), destroy the bone marrow and stem cells. After the high-dose treatment, you will be given a drip (infusion) of either your own stem cells (autologous transplant) or stem cells from a donor (allogeneic transplant).
- **Biological therapy** – also known as immunotherapy. This is the use of substances that are naturally produced within the body to encourage the immune system to fight disease. Interferon and growth factors are used to treat leukaemia.
- **Radiotherapy** – may be used for different types of leukaemia to help control symptoms as part of intensive treatment regimes, or to treat cancer that has spread to the brain or spinal cord.
- **Steroids** – research has shown that treatment for some types of leukaemia is better if steroids are given alongside the chemotherapy.
- **All-trans-retinoic acid** – this is a form of vitamin A used to treat a type of acute myeloid leukaemia called acute promyeloid leukaemia or APL. It is usually taken as tablets with chemotherapy.
- **Complementary and alternative therapies** – when used alongside your conventional cancer treatment, some of these therapies can make you feel better and improve quality of life. Others may not be so helpful and in some cases may be harmful. The Cancer Council Victoria booklet called **Understanding complementary therapies (pdf)** can be a useful resource.

All treatments have side effects. These will vary depending on the type of treatment you are having. Many side effects are temporary, but some may be permanent. Your doctor will explain all the possible side effects before

your treatment begins.

Research into leukaemia

Early detection and better treatment have improved survival for people with leukaemia. Research for leukaemia is ongoing. The [**Cancer Research UK**](#) website has information about research into all types of leukaemia.

Clinical trials can test the effectiveness of promising new treatments or new ways of combining cancer treatments. Always discuss treatment options with your doctor. There may be a clinical trial to suit your type of leukaemia.

Your sexuality and leukaemia

Having leukaemia and its treatment can affect the way you feel about your body, who you are, your relationships, the way you express yourself sexually and your sexual feelings (your 'sexuality'). These changes can be very upsetting.

Your medical team should discuss these issues with you before and during your treatment. If you feel you would like to discuss things further, ask your doctor for a referral to a counsellor or call the Cancer Information and Support Service on 13 11 20.

If you have a partner, it helps to be as open as possible with them about how you are feeling. The Cancer Council Victoria booklet called *Sexuality, intimacy and cancer* may also be helpful to read.

Caring for someone with leukaemia

Caring for someone with leukaemia can be a difficult and emotional time. If you or someone you know is caring for someone with leukaemia, there is support available. The Cancer Council Victoria booklet called *Caring for someone with cancer* may also be helpful to read.

When a cure isn't possible

If leukaemia has been diagnosed in its later stages, the cancer may have spread to the point where a cure is no longer possible. Treatment then focuses on improving quality of life by relieving the symptoms (this is called 'palliative' treatment). Medication can be used to relieve pain, nausea and vomiting. The Cancer Council Victoria booklet called *Living with advanced cancer* may be helpful to read.

Where to get help

- Your doctor
- **Cancer Council Victoria** Tel. **13 11 20**
- **Multilingual Cancer Information Line**, Victoria Tel. **13 14 50**
- **Leukaemia Foundation** Tel. 1800 620 420 or **(03) 9949 5800**
- **Leukaemia Foundation** has a number of inspiring **personal patient stories**
- **Leukaemia Family Support Group** Tel. **(03) 9342 7888**

This page has been produced in consultation with and approved by:

Cancer Council Victoria

Content on this website is provided for information purposes only. Information about a therapy, service, product or treatment does not in any way endorse or support such therapy, service, product or treatment and is not intended to replace advice from your doctor or other registered health professional. The information and materials contained on this website are not intended to constitute a comprehensive guide concerning all aspects of the therapy, product or treatment described on the website. All users are urged to always seek advice from a registered health care professional for diagnosis and answers to their medical questions and to ascertain whether the particular therapy, service, product or treatment described on the website is suitable in their circumstances. The State of Victoria and the Department of Health & Human Services shall not bear any liability for reliance by any user on the materials contained on this website.

For the latest updates and more information, visit www.betterhealth.vic.gov.au

Copyright © 1999/2019 State of Victoria. Reproduced from the Better Health Channel (www.betterhealth.vic.gov.au) at no cost with permission of the Victorian Minister for Health. Unauthorised reproduction and other uses comprised in the copyright are prohibited without permission.