
Contraception - diaphragms

Summary

- Diaphragms act as a barrier method of contraception.
 - Other barrier methods include male and female condoms.
 - Diaphragms need to stay in the vagina for at least six hours after sex.
 - Diaphragms do not give protection from sexually transmissible infections (STIs). The best way to lessen the risk of STIs is to use a condom with all new sexual partners.
 - For a diaphragm to be effective, you need to use it correctly, every time.
-

A diaphragm acts as a barrier method of contraception. It fits inside a woman's vagina and covers the cervix (entrance to the uterus or womb) to stop sperm from meeting and fertilising an egg. Other barrier methods include male and female condoms.

What is a diaphragm?

A diaphragm is a soft, shallow cup made of silicone. In Australia the one-size diaphragm is sold as Caya®. It is used with a special gel (jelly-like material, sold as Caya Gel®).

How effective are diaphragms?

If used correctly every time you have sex, diaphragms are 86 per cent effective at preventing pregnancy. If used incorrectly, for example not inserting properly before sex or removing too soon after sex, they may only be 82 per cent effective.

What stops diaphragms from working?

A diaphragm might not work if it is:

- not inserted correctly
- used without the gel
- taken out less than six hours after sex
- torn
- past its use-by date.

How do I use a diaphragm?

To use a diaphragm, place a teaspoon of gel in the cup of the diaphragm, then put the diaphragm inside your vagina so it covers the cervix (opening to your uterus).

Insert the diaphragm up to two hours before having sex, and leave it in place for at least six hours after sex. Do not leave the diaphragm in place for longer than 24 hours.

Before you use it, you can ask a doctor or nurse to check that the diaphragm fits well and that you are inserting it correctly. Diaphragms are the wrong shape for about one in seven women. The doctor or nurse can provide advice about how to insert the diaphragm comfortably. Caya® also provides a [**video with instructions about how to use the diaphragm**](#).

How does a diaphragm work?

The diaphragm and gel work as a barrier to prevent sperm from passing through the cervix into the uterus. The sperm die off quickly in the vagina.

Where can I get a diaphragm?

You can buy a diaphragm from some pharmacies, family planning clinics and online.

What is good about diaphragms?

The benefits of diaphragms are:

- You can insert them hours before having sex.
- They have little or no side effects.
- You can buy them without a script.
- They are a choice for women who do not want to use hormonal contraception.

Are there any side effects from using a diaphragm?

Possible side effects for a small number of users can include:

- irritation of the vagina or penis
- increased risk of bladder infection.

Can diaphragms cause any serious health problems?

Rarely, use of a diaphragm might cause **toxic shock syndrome**.

When is a diaphragm not a good option?

A diaphragm might not be a good option if you:

- are prone to bladder infections
- gave birth six weeks ago or less – because the size of diaphragm required may change as the uterus returns to normal size
- are unable to feel if your cervix is correctly covered by the diaphragm.

Speak to a doctor or nurse for advice.

What if I forget to use my diaphragm?

You may need emergency contraception if you had sex without using your diaphragm and did not use other contraception (check with your doctor, nurse or pharmacist).

What happens if I get pregnant while I'm using a diaphragm?

It is safe to continue the pregnancy (and stop using the diaphragm) or to have an abortion.

Can I use a diaphragm after I've had a baby?

You should usually wait six weeks after having a baby to start using a diaphragm.

What else should I know about diaphragms?

Diaphragms:

- do not protect you from STIs
- should be checked carefully for holes
- usually last around two years.

Where to get help

- Pharmacist
- **Family Planning Victoria** – comprehensive sexual and reproductive health services for people of all ages Tel. **1800 013 952** or **(03) 9257 0100**
- **Family Planning Victoria Action Centre** – comprehensive sexual and reproductive health services for people of all ages, with an afternoon drop-in clinic for people under 25 years of age Tel. **(03) 9660 4700** or **1800 013 952**
- **1800 my options** (**1800 696 784**) can provide information on the phone about contraception services

betterhealth.vic.gov.au

- Your **GP (doctor)**
- Many community health services and some public hospitals will have a family planning clinic, a sexual health clinic or a women's health clinic providing contraception
- Private clinics offering abortion also offer contraceptive services
- **Melbourne Sexual Health Centre** Tel. **(03) 9341 6200** or **1800 032 017**

This page has been produced in consultation with and approved by:

Family Planning Victoria

Content on this website is provided for information purposes only. Information about a therapy, service, product or treatment does not in any way endorse or support such therapy, service, product or treatment and is not intended to replace advice from your doctor or other registered health professional. The information and materials contained on this website are not intended to constitute a comprehensive guide concerning all aspects of the therapy, product or treatment described on the website. All users are urged to always seek advice from a registered health care professional for diagnosis and answers to their medical questions and to ascertain whether the particular therapy, service, product or treatment described on the website is suitable in their circumstances. The State of Victoria and the Department of Health & Human Services shall not bear any liability for reliance by any user on the materials contained on this website.

For the latest updates and more information, visit www.betterhealth.vic.gov.au

Copyright © 1999/2020 State of Victoria. Reproduced from the Better Health Channel (www.betterhealth.vic.gov.au) at no cost with permission of the Victorian Minister for Health. Unauthorised reproduction and other uses comprised in the copyright are prohibited without permission.