

---

## Receptive language disorder

---

### Summary

- A child with receptive language disorder has difficulties with understanding what is said to them.
- The cause of receptive language disorder is unknown, but is thought to consist of a number of factors working in combination.
- Speech-language therapy is used to treat receptive language disorder.

---

A child with receptive language disorder has difficulties with understanding what is said to them. The symptoms vary between children but, generally, problems with language comprehension begin before the age of three years.

Children need to understand spoken language before they can use language to express themselves. In most cases, children with a receptive language problem also have an **expressive language disorder**, which means they have trouble using spoken language.

It is estimated that between three and five per cent of children have a receptive or expressive language disorder, or a mixture of both. Another name for receptive language disorder is language comprehension deficit. Speech-language therapy is used to treat receptive language disorder.

### Symptoms of receptive language disorder

- There is no standard set of symptoms that indicates receptive language disorder, as it varies from one child to the next. However, symptoms may include:
  - not seeming to listen when they are spoken to
  - appearing to lack interest when storybooks are read to them
  - difficulty understanding the meaning of words and sentences
  - difficulty remembering all the words in a sentence in order to make sense of what has been said
  - inability to understand complicated sentences
  - inability to follow verbal instructions; especially if the instruction is long or complicated.

### Cause of receptive language disorder

The cause of receptive language disorder is often unknown, but is thought to consist of a number of factors working in combination, such as:

- genetic susceptibility (family history of receptive language disorder)
- limited exposure to hearing language in their day-to-day environment
- general developmental and cognitive (thinking) abilities.

Receptive language disorder is often associated with developmental disorders such as autism or Down syndrome. (Although for some children, difficulty with language is the only developmental problem they experience.)

In other cases, receptive language disorder is caused by damage to the brain, for example due to trauma, tumour or disease.

Receptive language disorder may also be related to:

- hearing impairment – due to decreased exposure to language
- vision impairment – due to the absence of cues such as facial expression and gestures

- attention disorders – due to difficulties in attending fully to what is being said.

### **Diagnosis of receptive language disorder**

Assessment needs to pinpoint the child's particular areas of difficulty, especially when they do not respond to spoken language. Diagnosis may include:

- hearing tests (by an audiologist) to check whether the language problems are caused by hearing impairment and to establish whether or not the child is able to pay attention to sound and language (auditory processing assessment)
- testing the child's language comprehension (by a speech pathologist) and comparing the results to the expected skill level for the child's age. If the child is from a non-English speaking home, assessment of comprehension should be performed in their first language, as well as in English, using culturally-appropriate materials
- close observation of the child in a variety of different settings while they interact with a range of people
- assessment by a psychologist to help identify any associated cognitive problems
- vision tests to check for vision impairment.

### **Treatment for receptive language disorder**

Treatment options for receptive language disorder may include:

- speech-language therapy (one-on-one or as part of a group, or both, depending on the needs of the child)
- providing information to families so that they can facilitate language growth at home
- special education classes at school
- integration support at preschool or school in cases of severe difficulty
- referral to a psychologist for treatment (only if there are also significant behavioural problems).

A child's progress will depend on a range of individual factors, such as whether or not brain injury is present.

### **Symptoms of expressive language disorder**

A child with receptive language disorder may also have expressive language disorder, which means they have difficulties with using spoken language. Symptoms differ from one child to the next, but can include:

- frequently grasping for the right word
- using the wrong words in speech
- making grammatical mistakes
- relying on short, simple sentence construction
- relying on stock standard phrases
- inability to 'come to the point' of what they're trying to say
- problems with retelling a story or relaying information
- inability to start or hold a conversation.

### **Where to get help**

- Your GP
- **Speech pathologist**
- **Audiologist**
- **Speech Pathology Australia** Tel. **(03) 9642 4899**
- Your maternal and child health nurse.

**This page has been produced in consultation with and approved by:**

La Trobe University - School of Allied Health - Speech Pathology

Content on this website is provided for information purposes only. Information about a therapy, service, product or treatment does not in any way endorse or support such therapy, service, product or treatment and is not intended to replace advice from your doctor or other registered health professional. The information and materials contained on this website are not intended to constitute a comprehensive guide concerning all aspects of the therapy, product or treatment described on the website. All users are urged to always seek advice from a registered health care professional for diagnosis and answers to their medical questions and to ascertain whether the particular therapy, service, product or treatment described on the website is suitable in their circumstances. The State of Victoria and the Department of Health & Human Services shall not bear any liability for reliance by any user on the materials contained on this website.

For the latest updates and more information, visit [www.betterhealth.vic.gov.au](http://www.betterhealth.vic.gov.au)

**Copyright © 1999/2020** State of Victoria. Reproduced from the Better Health Channel ([www.betterhealth.vic.gov.au](http://www.betterhealth.vic.gov.au)) at no cost with permission of the Victorian Minister for Health. Unauthorised reproduction and other uses comprised in the copyright are prohibited without permission.